

The geometry of fluvial channel bodies: Empirical characterization and implications for object-based models of the subsurface

Luca Colombera, Nigel P. Mountney, Giacomo Medici, and L. Jared West

AAPG Bulletin, v. 103, no. 4 (April 2019), pp. 905–929

Copyright ©2019. The American Association of Petroleum Geologists. All rights reserved.

LIST OF PUBLISHED SOURCES THAT CONTAIN DATA USED IN THIS WORK

- Alqahtani, F. A., C. A.-L. Jackson, H. D. Johnson, and M. R. B. Som, 2017, Controls on the geometry and evolution of humid-tropical fluvial systems: Insights from 3D seismic geomorphological analysis of the Malay Basin, Sunda Shelf, Southeast Asia: *Journal of Sedimentary Research*, v. 87, no. 1, p. 17–40, doi:[10.2110/jsr.2016.88](https://doi.org/10.2110/jsr.2016.88).
- Alqahtani, F. A., H. D. Johnson, C. A.-L. Jackson, and M. R. B. Som, 2015, Nature, origin and evolution of a Late Pleistocene incised valley-fill, Sunda Shelf, Southeast Asia: *Sedimentology*, v. 62, no. 4, p. 1198–1232, doi:[10.1111/sed.12185](https://doi.org/10.1111/sed.12185).
- Amorosi, A., M. Pavese, M. Ricci Lucchi, G. Sarti, and A. Piccin, 2008, Climatic signature of cyclic fluvial architecture from the Quaternary of the central Po Plain, Italy: *Sedimentary Geology*, v. 209, no. 1–4, p. 58–68, doi:[10.1016/j.sedgeo.2008.06.010](https://doi.org/10.1016/j.sedgeo.2008.06.010).
- Berendsen, H. J. A., and E. Stouthamer, 2001, Palaeogeographic development of the Rhine-Meuse delta, The Netherlands: Assen, the Netherlands, Koninklijke Van Gorcum, 268 p., doi:[10.1046/j.1365-3091.2002.00469_1.x](https://doi.org/10.1046/j.1365-3091.2002.00469_1.x).
- Bridge, J. S., G. A. Jalfin, and S. M. Georgieff, 2000, Geometry, lithofacies, and spatial distribution of Cretaceous fluvial sandstone bodies, San Jorge Basin, Argentina: Outcrop analog for the hydrocarbon-bearing Chubut Group: *Journal of Sedimentary Research*, v. 70, no. 2, p. 341–359, doi:[10.1306/2DC40915-0E47-11D7-8643000102C1865D](https://doi.org/10.1306/2DC40915-0E47-11D7-8643000102C1865D).
- Carter, D. C., 2003, 3-D seismic geomorphology: Insights into fluvial reservoir deposition and performance, Widuri field, Java Sea: *AAPG Bulletin*, v. 87, no. 6, p. 909–934, doi:[10.1306/01300300183](https://doi.org/10.1306/01300300183).
- Cuevas Martínez, J. L., L. Cabrera Pérez, A. Marcuello, P. Arbués Cazo, M. Marzo Carpio, and F. Bellmunt, 2010, Exhumed channel sandstone networks within fluvial fan deposits from the Oligo-Miocene Caspe Formation, South-east Ebro Basin (North-east Spain): *Sedimentology*, v. 57, p. 162–189, doi:[10.1111/j.1365-3091.2009.01096.x](https://doi.org/10.1111/j.1365-3091.2009.01096.x).
- Dalrymple, M., 2001, Fluvial reservoir architecture in the Statfjord Formation (northern North Sea) augmented by outcrop analogue statistics: *Petroleum Geoscience*, v. 7, no. 2, p. 115–122, doi:[10.1144/petgeo.7.2.115](https://doi.org/10.1144/petgeo.7.2.115).
- Darmadi, Y., B. J. Willis, and S. L. Dorobek, 2007, Three-dimensional seismic architecture of fluvial sequences on the low-gradient Sunda Shelf, offshore Indonesia: *Journal of Sedimentary Research*, v. 77, no. 3, p. 225–238, doi:[10.2110/jsr.2007.024](https://doi.org/10.2110/jsr.2007.024).
- Donselaar, M. E., and I. Overeem, 2008, Connectivity of fluvial point-bar deposits: An example from the Miocene Huesca fluvial fan, Ebro Basin, Spain: *AAPG Bulletin*, v. 92, no. 9, p. 1109–1129, doi:[10.1306/04180807079](https://doi.org/10.1306/04180807079).
- Dreyer, T., 1993, Quantified fluvial architecture in ephemeral stream deposits of the Esplugafreda Formation (Palaeanocene), Tremp-Graus basin, northern Spain, in M. Marzo and C. Puidefabregas, eds., *Alluvial sedimentation*: Hoboken, New Jersey, International Association of Sedimentologists Special Publication 17, p. 337–362, doi:[10.1002/9781444303995.ch23](https://doi.org/10.1002/9781444303995.ch23).
- Erhardt, A. M., 2005, Relative contributions of tectonics and climate on fluvial sedimentation in the Wasatch Formation of western Colorado, M.Sc. thesis, Colorado School of Mines, Golden, Colorado, 118 p.
- Fachmi, M., and L. J. Wood, 2005, Seismic geomorphology: A study from West Natuna Basin, Indonesia: 30th Annual Indonesian Petroleum Association Convention, Jakarta, Indonesia, August 30–September 1, 2005, p. 163–178.
- Fielding, C. R., 1986, Fluvial channel and overbank deposits from the Westphalian of the Durham coalfield, NE England.: *Sedimentology*, v. 33, no. 1, p. 119–140, doi:[10.1111/j.1365-3091.1986.tb00748.x](https://doi.org/10.1111/j.1365-3091.1986.tb00748.x).
- Fielding, C. R., and R. C. Crane, 1987, An application of statistical modelling to the prediction of hydrocarbon recovery factors in fluvial reservoir sequences, in F. G. Ethridge, R. M. Flores, and M. D. Harvey, eds., *Recent developments in fluvial sedimentology*: Tulsa, Oklahoma, SEPM Special Publication 39, p. 321–327, doi:[10.2110/pec.87.39.0321](https://doi.org/10.2110/pec.87.39.0321).
- Flores, R. M., and G. D. Stricker, 1993, Early Cenozoic depositional systems, Wishbone Hill District, Matanuska coal field, Alaska, in C. Dusel-Bacon and A. B. Till, eds., *Geologic studies in Alaska by the Geological Survey, 1992*: Washington, DC, US Geological Survey Bulletin 2068, p. 101–117.
- Ford, G. L., and D. R. Pyles, 2014, A hierarchical approach for evaluating fluvial systems: Architectural analysis and sequential evolution of the high net-sand content, middle Wasatch Formation, Uinta Basin, Utah: *AAPG Bulletin*, v. 98, no. 7, p. 1273–1304, doi:[10.1306/12171313052](https://doi.org/10.1306/12171313052).
- Foreman, B. Z., 2014, Climate-driven generation of a fluvial sheet sand body at the Paleocene–Eocene boundary in north-west Wyoming (USA): *Basin Research*, v. 26, no. 2, p. 225–241, doi:[10.1111/bre.12027](https://doi.org/10.1111/bre.12027).
- Foreman, B. Z., P. L. Heller, and M. T. Clementz, 2012, Fluvial response to abrupt global warming at the

- Palaeocene/Eocene boundary: *Nature*, v. 491, no. 7422, p. 92–95, doi:[10.1038/nature11513](https://doi.org/10.1038/nature11513).
- Friend, P. F., S. M. Raza, G. Geehan, and K. A. Sheikh, 2001, Intermediate-scale architectural features of the fluvial Chinji Formation (Miocene), Siwalik Group, northern Pakistan: *Journal of the Geological Society*, v. 158, p. 163–177, doi:[10.1144/jgs.158.1.163](https://doi.org/10.1144/jgs.158.1.163).
- Gouw, M. J. P., 2008, Alluvial architecture of the Holocene Rhine–Meuse delta (the Netherlands): *Sedimentology*, v. 55, no. 5, p. 1487–1516, doi:[10.1111/j.1365-3091.2008.00954.x](https://doi.org/10.1111/j.1365-3091.2008.00954.x).
- Gouw, M. J. P., and H. J. A. Berendsen, 2007, Variability of channel-belt dimensions and the consequences for alluvial architecture: Observations from the Holocene Rhine–Meuse Delta (the Netherlands) and Lower Mississippi Valley (USA): *Journal of Sedimentary Research*, v. 77, no. 2, p. 124–138, doi:[10.2110/jsr.2007.013](https://doi.org/10.2110/jsr.2007.013).
- Hajek, E. A., P. L. Heller, and B. A. Sheets, 2010, Significance of channel-belt clustering in alluvial basins: *Geology*, v. 38, no. 6, p. 535–538, doi:[10.1130/G30783.1](https://doi.org/10.1130/G30783.1).
- Hampson, G. J., M. R. Gani, H. Sahoo, A. Rittersbacher, N. Irfan, A. Ranson, T. O. Jewell et al., 2012, Controls on large-scale patterns of fluvial sandbody distribution in alluvial to coastal plain strata: Upper Cretaceous Blackhawk Formation, Wasatch Plateau, Central Utah, USA: *Sedimentology*, v. 59, no. 7, p. 2226–2258, doi:[10.1111/j.1365-3091.2012.01342.x](https://doi.org/10.1111/j.1365-3091.2012.01342.x).
- Hirst, J. P. P., 1991, Variations in alluvial architecture across the Oligo-Miocene Huesca fluvial system, Ebro Basin, Spain, in A. D. Miall and N. Tyler, eds., *The three-dimensional facies architecture of terrigenous clastic sediments and its implications for hydrocarbon discovery and recovery*: Tulsa, Oklahoma, SEPM Concepts in Sedimentology and Paleontology 3, p. 111–121.
- Jo, H. R., 2003, Depositional environments, architecture, and controls of Early Cretaceous non-marine successions in the northwestern part of Kyongsang Basin, Korea: *Sedimentary Geology*, v. 161, no. 3–4, p. 269–294, doi:[10.1016/S0037-0738\(03\)00130-1](https://doi.org/10.1016/S0037-0738(03)00130-1).
- Jordan, D. W., and W. A. Pryor, 1992, Hierarchical levels of heterogeneity in a Mississippi River meander belt and application to reservoir systems: *AAPG Bulletin*, v. 76, no. 10, p. 1601–1624.
- Keighley, D., S. Flint, J. Howell, and A. Moscariello, 2003, Sequence stratigraphy in lacustrine basins: A model for part of the Green River Formation (Eocene), southwest Uinta Basin, Utah, U.S.A.: *Journal of Sedimentary Research*, v. 73, no. 6, p. 987–1006, doi:[10.1306/050103730987](https://doi.org/10.1306/050103730987).
- Kjemperud, A. V., E. R. Schomacker, and T. A. Cross, 2008, Architecture and stratigraphy of alluvial deposits, Morrison Formation (Upper Jurassic), Utah: *AAPG Bulletin*, v. 92, no. 8, p. 1055–1076, doi:[10.1306/03250807115](https://doi.org/10.1306/03250807115).
- Kraus, M. J., 1996, Avulsion deposits in lower Eocene alluvial rocks, Bighorn Basin, Wyoming: *Journal of Sedimentary Research*, v. 66, p. 354–363.
- Kraus, M. J., and A. Aslan, 1993, Eocene hydromorphic paleosols: Significance for interpreting ancient floodplain processes: *Journal of Sedimentary Research*, v. 63, p. 453–463, doi:[10.1306/D4267B22-2B26-11D7-8648000102C1865D](https://doi.org/10.1306/D4267B22-2B26-11D7-8648000102C1865D).
- Kraus, M. J., and K. S. Davies-Vollum, 2004, Mudrock-dominated fills formed in avulsion splay channels: Examples from the Willwood Formation, Wyoming: *Sedimentology*, v. 51, no. 5, p. 1127–1144, doi:[10.1111/j.1365-3091.2004.00664.x](https://doi.org/10.1111/j.1365-3091.2004.00664.x).
- Kraus, M. J., and B. Gwinn, 1997, Facies and facies architecture of paleogene floodplain deposits, Willwood formation, Bighorn Basin, Wyoming, USA: *Sedimentary Geology*, v. 114, no. 1–4, p. 33–54, doi:[10.1016/S0037-0738\(97\)00083-3](https://doi.org/10.1016/S0037-0738(97)00083-3).
- Labourdette, R., 2011, Stratigraphy and static connectivity of braided fluvial deposits of the lower Escanilla Formation, south central Pyrenees, Spain: *AAPG Bulletin*, v. 95, no. 4, p. 585–617, doi:[10.1306/08181009203](https://doi.org/10.1306/08181009203).
- Makaske, B., H. J. A. Berendsen, and M. H. M. van Ree, 2007, Middle Holocene avulsion-belt deposits in the Central Rhine–Meuse Delta, The Netherlands: *Journal of Sedimentary Research*, v. 77, no. 2, p. 110–123, doi:[10.2110/jsr.2007.004](https://doi.org/10.2110/jsr.2007.004).
- Martinius, A. W., and R. A. Nieuwenhuijs, 1995, Geological description of flow units in channel sandstones in a fluvial reservoir analogue (Loranca Basin, Spain): *Petroleum Geoscience*, v. 1, no. 3, p. 237–252, doi:[10.1144/petgeo.1.3.237](https://doi.org/10.1144/petgeo.1.3.237).
- Maynard, J. R., 2006, Fluvial response to active extension: Evidence from 3D seismic data from the Frio Formation (Oligo-Miocene) of the Texas Gulf of Mexico Coast, USA: *Sedimentology*, v. 53, no. 3, p. 515–536, doi:[10.1111/j.1365-3091.2006.00782.x](https://doi.org/10.1111/j.1365-3091.2006.00782.x).
- Maynard, J. R., H. R. Feldman, and R. Alway, 2010, From bars to valleys: The sedimentology and seismic geomorphology of fluvial to estuarine incised-valley fills of the Grand Rapids Formation (Lower Cretaceous), Iron River Field, Alberta, Canada: *Journal of Sedimentary Research*, v. 80, no. 7, p. 611–638, doi:[10.2110/jsr.2010.060](https://doi.org/10.2110/jsr.2010.060).
- Maynard, K., and I. Murray, 2003, One million years from the Upper Arang Formation, West Natuna Basin, implications for reservoir distribution and facies variation in fluvial deltaic deposits: 29th Annual Indonesian Petroleum Association Convention, Jakarta, Indonesia, October 14–16, 2003, p. 270–276.
- Maynard, K., P. Siregar, and L. Andria, 2002, Seismic stratigraphic interpretation of a major 3D, the Gabus sub-basin, blocks B and Tobong, West Natuna Sea, Indonesia: Getting the geology back into seismic: 28th Annual Indonesian Petroleum Association Convention, Jakarta, Indonesia, February 26–28, 2002, p. 87–104.
- McRae, L. E., 1990, Paleomagnetic isochrons, unsteadiness, and non-uniformity of sedimentation in Miocene fluvial strata of the Siwalik Group, Northern Pakistan: *Journal of Geology*, v. 98, no. 4, p. 433–456, doi:[10.1086/629418](https://doi.org/10.1086/629418).
- Meadows, N. S., 2006, The correlation and sequence architecture of the Ormskirk Sandstone Formation in the Triassic Sherwood Sandstone Group of the East Irish Sea Basin, NW England: *Geological Journal*, v. 41, no. 1, p. 93–122, doi:[10.1002/gj.1034](https://doi.org/10.1002/gj.1034).
- Miall, A. D., 2002, Architecture and sequence stratigraphy of Pleistocene fluvial systems in the Malay Basin, based on seismic time-slice analysis: *AAPG Bulletin*, v. 86,

- no. 7, p. 1201–1216, doi:[10.1306/61EEDC56-173E-11D7-8645000102C1865D](https://doi.org/10.1306/61EEDC56-173E-11D7-8645000102C1865D).
- Mohrig, D., P. L. Heller, C. Paola, and W. J. Lyons, 2000, Interpreting avulsion process from ancient alluvial sequences: Guadalupe-Matarranya system (northern Spain) and Wasatch Formation (western Colorado): Geological Society of America Bulletin, v. 112, no. 12, p. 1787–1803, doi:[10.1130/0016-7606\(2000\)112<1787:IAPFAA>2.0.CO;2](https://doi.org/10.1130/0016-7606(2000)112<1787:IAPFAA>2.0.CO;2).
- Nádor, A., and O. Zstanó, 2011, Lateral and vertical variability of channel belt stacking density as a function of subsidence and sediment supply: Field evidence from the intra-mountain Körös Basin, Hungary, in S. K. Davidson, S. Leleu, and C. P. North, eds., From river to rock record: The preservation of fluvial sediments and their subsequent interpretation: Tulsa, Oklahoma, SEPM Special Publication 97, p. 375–392, doi:[10.2110/sepmfsp.097.375](https://doi.org/10.2110/sepmfsp.097.375).
- North, C. P., and K. S. Taylor, 1996, Ephemeral-fluvial deposits: Integrated outcrop and simulation studies reveal complexity: AAPG Bulletin, v. 80, no. 6, p. 811–830.
- Olsen, T., 1995a, Fluvial and fluvio-lacustrine facies and depositional environments of the Maastrichtian to Paleocene North Horn Formation, Price Canyon, Utah: The Mountain Geologist, v. 32, p. 27–44.
- Olsen, T., 1995b, Sequence stratigraphy, alluvial architecture and potential reservoir heterogeneities of fluvial deposits: Evidence from outcrop studies in Price Canyon, Utah (Upper Cretaceous and Lower Tertiary), in R. J. Steel, V. L. Felt, E. P. Johannessen, C. Mathieu, eds., Sequence stratigraphy on the northwest European margin: Amsterdam, Norwegian Petroleum Society Special Publication 5, p. 75–88, 92–96, doi:[10.1016/S0928-8937\(06\)80064-0](https://doi.org/10.1016/S0928-8937(06)80064-0).
- Paredes, J. M., N. Foix, F. Colombo Piñol, A. Nillni, J. O. Allard, and R. A. Marquillas, 2007, Volcanic and climatic controls on fluvial style in a high-energy system: The Lower Cretaceous Matasiete Formation, Golfo San Jorge basin, Argentina: Sedimentary Geology, v. 202, no. 1–2, p. 96–123, doi:[10.1016/j.sedgeo.2007.05.007](https://doi.org/10.1016/j.sedgeo.2007.05.007).
- Rasmussen, A. M. S., 2005, Reservoir characterization of a fluvial sandstone: Depositional environment and heterogeneities in modeling of the Colton Formation, Utah, M.Sc. thesis, University of Oslo, Oslo, Norway, 136 p.
- Reijenstein, H. M., H. W. Posamentier, and J. P. Bhattacharya, 2011, Seismic geomorphology and high-resolution seismic stratigraphy of inner-shelf fluvial, estuarine, deltaic, and marine sequences, Gulf of Thailand: AAPG Bulletin, v. 95, no. 11, p. 1959–1990, doi:[10.1306/03151110134](https://doi.org/10.1306/03151110134).
- Reynolds, A. D., 1999, Dimensions of paralic sandstone bodies: AAPG Bulletin, v. 83, no. 2, p. 211–229, doi:[10.1306/00AA9A48-1730-11D7-8645000102C1865D](https://doi.org/10.1306/00AA9A48-1730-11D7-8645000102C1865D).
- Roberts, E. M., 2007, Facies architecture and depositional environments of the Upper Cretaceous Kaiparowits Formation, southern Utah: Sedimentary Geology, v. 197, no. 3–4, p. 207–233, doi:[10.1016/j.sedgeo.2006.10.001](https://doi.org/10.1016/j.sedgeo.2006.10.001).
- Roberts, S. B., 1998, An overview of the stratigraphic and sedimentologic characteristics of the Paleocene Fort Union Formation, southern Bighorn Basin, Wyoming, in W. R. Keefer and J. E. Goolsby, eds., Cretaceous and lower Tertiary rocks of the Bighorn Basin, Wyoming and Montana: Wyoming Geological Association Guidebook 49, p. 91–116.
- Robinson, J. W., and P. J. McCabe, 1997, Sandstone-body and shale-body dimensions in a braided fluvial system: Salt Wash Sandstone Member (Morrison Formation), Garfield County, Utah: AAPG Bulletin, v. 81, no. 8, p. 1267–1291.
- Rogers, R. R., 1998, Sequence analysis of the Upper Cretaceous Two Medicine and Judith River formations, Montana: Nonmarine response to the Claggett and Bearpaw marine cycles: Journal of Sedimentary Research, v. 68, no. 4, p. 615–631, doi:[10.2110/jsr.68.604](https://doi.org/10.2110/jsr.68.604).
- Rygel, M. C., and M. R. Gibling, 2006, Natural geomorphic variability recorded in a high-accommodation setting: Fluvial architecture of the Pennsylvanian Joggins Formation of Atlantic Canada: Journal of Sedimentary Research, v. 76, no. 11, p. 1230–1251, doi:[10.2110/jsr.2006.100](https://doi.org/10.2110/jsr.2006.100).
- Samorn, H., 2006, Fluvial reservoir architecture from near-surface 3D seismic data, block B8/32, Gulf of Thailand, M.Sc. thesis, Colorado School of Mines, Golden, Colorado, 156 p.

ADDITIONAL DATA WERE DERIVED FROM OUTCROP STUDIES. PUBLICATIONS THAT CONTAIN THE SAME DATA SETS ARE REPORTED BELOW

- Colombera, L., O. J. Arévalo, and N. P. Mountney, 2017, Fluvial-system response to climate change: The Paleocene-Eocene Tremp Group, Pyrenees, Spain: Global and Planetary Change, v. 157, p. 1–17, doi:[10.1016/j.gloplacha.2017.08.011](https://doi.org/10.1016/j.gloplacha.2017.08.011).
- Colombera, L., N. P. Mountney, J. A. Howell, A. Rittersbacher, F. Felletti, and W. D. McCaffrey, 2016, A test of analog-based tools for quantitative prediction of large-scale fluvial architecture: AAPG Bulletin, v. 100, no. 2, p. 237–267, doi:[10.1306/11181514227](https://doi.org/10.1306/11181514227).